

On a question of Vaisman concerning complex surfaces

Massimiliano Pontecorvo

May, 2012

In memory of Marco Brunella

Abstract

The last years have seen striking improvements on Vaisman's question about existence of locally conformally Kähler (lcK) metrics on compact complex surfaces. The aim of this paper is two-fold. We review results of different authors which, for all known examples of compact complex surfaces, give a complete answer to Vaisman's question. We also point out a relation between lcK surfaces and generalized Kähler geometry in four-dimension and prove a new result concerning generalized Kähler structures on Hyperbolic Inoue surfaces. We conclude with a simple observation on a question of Brunella.

1 Introduction

Throughout this work M will denote a smooth oriented 4-manifold - which will be compact most of the times and always connected - equipped with a Riemannian metric g , but in fact most of our notions will only depend on the conformal class $[g]$ of all metrics $g' = e^f g$ which are obtained multiplying g by a positive smooth function.

(M, g, J) is said to be a Hermitian surface when J is a g -orthogonal endomorphism of the tangent bundle $J : TM \rightarrow TM$ with $J^2 = -id$. inducing holomorphic coordinates, so that (M, J) is a smooth complex manifold and $\dim_{\mathbb{C}} M = 2$. Because J is orthogonal for any metric in $[g]$, we will often refer to the Hermitian conformal surface $(M, [g], J)$.

The *Hermitian 2-form* of the metric (g, J) is defined to be $\Omega(\cdot, \cdot) := g(\cdot, J\cdot)$ and in four-dimensions defines an isomorphism between 1-forms and 3-forms so that

$$d\Omega = \Omega \wedge \theta$$

for a unique 1-form θ called the *Lee form* of the Hermitian metric; since the metric is Kähler if and only if $\theta = 0$ and the Lee form θ' of a conformally related metric $g' = e^f g$ is given by $\theta' = \theta + df$, we easily have:

1. The Lee form $\theta = df$ is exact when the metric $e^f g$ is Kähler - i.e. (g, J) is a globally conformally Kähler metric.
2. The Lee form is closed - i.e. locally exact, $d\theta = 0$ - precisely when the Hermitian metric is conformal to Kähler, locally.

Such metrics are then called *locally conformally Kähler*, usually abbreviated by lcK. Of course, it is a conformally invariant notion.

The paper is dedicated to discuss the following question of Vaisman:

Question 1.1 [Va87, p.122] *Which compact complex surfaces (M, J) admit lcK metrics?*

By a celebrated result of Gauduchon, in the compact four-dimensional case, there always exists a unique metric in the conformal class - called the Gauduchon metric of $(M, [g], J)$ - whose Lee form θ is coclosed. Furthermore θ is coexact if and only if the first Betti number $b_1(M)$ of M is even [Ga84].

As a consequence, when $b_1(M)$ is even, the Lee form of a Gauduchon metric on (M, J) can be closed only for Kähler metrics. It follows that lcK compact surfaces with b_1 even are automatically globally conformally Kähler; this had already been proved by Vaisman in a different way [Va80]. The conclusion is that, for $b_1(M)$ even, Vaisman question coincides with a famous conjecture of Kodaira:

Conjecture 1.2 [KM71] *A compact complex surface is Kählerian if and only if the first Betti number is even.*

This conjecture became a Theorem since the early 1980's and was first proved using the Enriques-Kodaira classification of surfaces with the contribution of Miyaoka, Todorov and Siu who clarified [Si83] the remaining case of K3 surfaces. It is also interesting to notice that two independent, direct proofs by Buchdahl and Lamari - not using surface classification - have simultaneously appeared more recently [Bu99], [La99].

Because any compact complex surface can be obtained by blowing up points on a minimal model, the Enriques-Kodaira classification only takes care of *minimal* surfaces - i.e. without smooth rational curves of self-intersection (-1) - and fits very well with the following result of Tricerri which generalizes the analogous well known result of Kodaira in the Kähler case; see also the work of Vuletescu for a more recent reference:

Proposition 1.3 [Tr82], [Vu09] *A complex manifold M is lcK if and only if the blow up of M at a point is lcK*

As noticed in [Tr82, Remark 4.3] this reduces Vaisman question to minimal surfaces and it is quite natural to use the Enriques-Kodaira classification to attack the problem.

For the reader's convenience we will display - at the end of this section - a (slightly refined) table [BHPV04, p.244] of this well known classification which divides all minimal compact complex surfaces $S := (M, J)$ in eleven classes belonging to four groups according to the possible values of the *Kodaira dimension* $Kod(S) = -\infty, 0, 1, 2$, which appears in the second column of the table; in the other columns are indicated: the *algebraic dimension* $a(S)$, which is the degree of the extension (over $\mathbb{C}$) of the field of meromorphic functions on M , the Euler characteristic $\chi(S) = 2 - 2b_1(S) + b_2(S)$ and the first Betti number $b_1(S)$.

We believe that Vaisman question is a good one because of its intricate relations with the study of complex geometry of non-Kähler surfaces developed by the Japanese school of Kodaira, Nakamura, Inoue, Kato, Enoki and more recently by the Marseille-based group of Dloussky, Teleman, Oeljeklaus, Toma.

The remaining of the paper is divided in three sections; the first is dedicated to the work of Belgun [Be00] who gave new impetus to the conjecture by completing the work initiated by Vaisman, Tricerri and Gauduchon-Ornea [GO99] on complex surfaces of vanishing Euler characteristic. The case of class-VII surfaces appears to be the most difficult and this parallels the fact that the classification of complex structures on these surfaces ($\chi = 0$) was solved by Bogomolov only in 1976 [Bg76]. The proof was later clarified by Li-Yau-Zheng [LYZ94] and Teleman [Te94], independently.

One major discovery of Belgun is that - unlike in the Kähler case - lcK is not an open property under holomorphic deformations. More precisely, he showed that some of the Inoue-Bombieri surfaces [In74], [Bm73] are not lcK. As a matter of fact this Hermitian property is somehow reflected in the complex geometry: Inoue-Bombieri surfaces are the only known compact complex surfaces with no curves.

After the work of Belgun, Vaisman question had a complete (although not always positive) answer for all surfaces except for those with minimal model satisfying $b_1 = 1$ and $b_2 > 0$; they are usually said to belong to class VII_0^+ . In section 3, we present our contribution about Inoue surfaces in this class.

It is important at this point to spend a few words on the classification problem of class VII_0^+ surfaces, the best reference for known results and open questions are the expositions of Nakamura [Na89], [Na08]. It is not even known - for example - whether every $S \in VII_0^+$ has

a (holomorphic) curve; a remarkable, recent result of A. Teleman [Te10] implies that this is the case when $b_2(S) \leq 2$.

As a matter of fact, the only known examples in this class are so called Kato surfaces [Ka77]: they can be characterized by saying that they have $b_2(S)$ rational curves [DOT03]. Among them, the Inoue surfaces [In77] form a subclass in which every rational curve belongs to a cycle $C \subset S$. These Inoue surfaces were the first examples of lcK surfaces in class VII_0^+ . Their lcK metrics came from a link - pointed out by Boyer, see Prop. 3.1 - with self-duality in dimension four.

Surprisingly enough, it turns out that many Inoue surfaces in class VII_0^+ carry bi-Hermitian metrics. This provides a new relation between lcK and generalized Kähler geometry [Gu07], [Hi06] given by the fact that anti-self-dual bi-Hermitian metrics always satisfy the generalized Kähler property that the sum of the two Lee forms vanishes identically [Po97, 3.11].

We present a new result in this context; Theorem 3.6 proves existence of generalized Kähler metrics on all admissible blow-ups of Hyperbolic Inoue surfaces. The first lcK examples in class VII_0^+ are by LeBrun [Le91] on Parabolic Inoue surfaces; later, a twistor construction by Fujiki-Pontecorvo [FP10] produced these very special lcK metrics on all (minimal) Hyperbolic as well as on all Half-Inoue surfaces.

Shortly after came the remarkable contribution of Marco Brunella [Br10], [Br11], which is the subject of the last section. He proved that - contrary to the case of Inoue-Bombieri surfaces - the lcK property is open under holomorphic deformations of Kato surfaces. In fact, he showed furthermore that all Kato surfaces are lcK.

Once again, Brunella results reflect the classification problem of non-Kähler surfaces: indeed, Kato surfaces are the only known surfaces in class VII_0^+ .

Vaisman question would now have a complete answer if no other class- VII_0^+ surfaces would exist. This is in fact a strong conjecture of Nakamura [Na89, 5.5].

Much more recently A. Teleman, using Gauge theory, has proved this conjecture for $b_2 = 1$ in [Te05]; he also proved in [Te10] that for $b_2 = 2$ every class- VII_0^+ surface contains a cycle of rational curves.

We conclude the paper with a simple observation on a question of Brunella [Br11].

Acknowledgement. A preliminary and incomplete version of this work appeared in the informal volume on the occasion of Fujiki's 60th birthday [Po08]. We take this opportunity to thank the organizers of the conference.

Table of Enriques-Kodaira classification

Class of S	$Kod(S)$	$a(S)$	$\chi(S)$	$b_1(S)$
1) rational surfaces 2) ruled surfaces of genus $g \geq 1$ 3) class VII_0 surfaces 4) class VII_0^+ surfaces	$-\infty$	2 2 0,1 0	3,4 $4(1-g)$ 0 > 0	0 $2g$ 1 1
5) tori 6) K3-surfaces 7) hyperelliptic surfaces 8) Enriques surfaces 9) Kodaira surfaces	0	0,1,2 0,1,2 2 2 1	0 24 0 12 0	4 0 2 0 1,3
10) properly elliptic surfaces	1	1,2 1	≥ 0 0	even odd
11) surfaces of general type	2	2	> 0	even

2 Non-Kähler surfaces with $\chi = 0$

After reducing the problem to minimal surfaces with $b_1(S)$ odd, we see from Kodaira classification that the Euler characteristic satisfies $\chi(S) \geq 0$. Extending previous results of Vaisman [Va87], Tricerri [Tr82] and Gauduchon-Ornea [GO99], Belgun gave a complete answer to Vaisman question in the limiting case when $\chi(S) = 0$, see [Be00]. In fact, because $\chi(S) = 0$ is clearly a necessary condition for a lcK metric to have parallel Lee form, Belgun even classified compact complex surfaces with these special lcK metrics which are called Vaisman metrics. The main tool was the use of explicit models for the universal cover of such geometric complex surfaces, which we now describe in order of decreasing Kodaira dimension. Recall that in what follows we assume $b_1(S)$ odd and $\chi(S) = 0$ which implies S minimal, by Kodaira classification and the fact that χ increases under blow up.

$$\underline{Kod(S) = 1}$$

These are so called properly elliptic surfaces which means that they are elliptic of algebraic dimension 1. In general, $Kod(S) \leq a(S)$ and in our case equality holds because every Mois-chezon surface is algebraic [BHPV04]. Then S has a holomorphic map to a curve with connected fibers and the generic fiber is elliptic, otherwise it would have non-zero intersection with the canonical class and S would be algebraic. Elliptic surfaces were studied in depth by Kodaira [Ko64] and when b_1 is odd, the only singular fibers turn out to be multiple fibers. Maheara [Ma77] showed that the universal cover is $\mathbb{C} \times H$ the product of a complex line with the upper half plane. The explicit description of the covering map allows Belgun to prove that all minimal surfaces with odd b_1 and $Kod(S) = 1$ admit lcK metrics and in fact Vaisman metrics.

$$\underline{Kod(S) = 0}$$

These are called Kodaira surfaces and are divided into primary or secondary according to whether b_1 is 1 or 3. Primary Kodaira surfaces $S = (M, J)$ are elliptic fiber bundles over an elliptic curve. It was shown by Thurston that they also admit a symplectic structure I and (M, I) provided the first example of a compact symplectic manifold without Kähler structures [Th76]. Finally, Salamon observed in [Sa94] that I and J anti commute and therefore give rise to an almost hyperHermitian structure spanned by $\{I, J, IJ\}$ with one complex structure J and two symplectic structures I and IJ .

A secondary Kodaira surface is finitely covered by a primary one and the universal covering is a nilpotent group; Belgun showed that all of these surfaces always admit Vaisman metrics, in particular lcK metrics.

$$\underline{Kod(S) = -\infty}$$

This is the class of VII_0 -surfaces in the above table, they are characterized by having negative Kodaira dimension and vanishing second Betti number; the already cited Bogomolov theorem [Bg76] states they are either Hopf surfaces - which always have at least one curve - or the surfaces independently discovered at the same time by Inoue [In74] and Bombieri [Bm73] which have no curves at all; we call them Inoue-Bombieri surfaces in order to distinguish from a different type of Inoue surfaces [In77] which belong to class VII_0^+ and will be the subject of next section.

Hopf surfaces have universal covering $\mathbb{C}^2 \setminus 0$ on which we take global coordinates (z, w) ; the primary Hopf surfaces are diffeomorphic to $S^1 \times S^3$ with fundamental group generated by the contraction

$$(z, w) \mapsto (az, bw + \lambda z^n)$$

where $a, b, \lambda \in \mathbb{C}$ and $n \in \mathbb{N}$ satisfy $0 < |a| < |b| < 1$ and $\lambda(a - b^n) = 0$; we say that a Hopf surface is *diagonal* when $\lambda = 0$ and in this case there are always (at least) two (elliptic) curves

on S , namely the images of the two axes $\{z = 0\}$ and $\{w = 0\}$. When $\lambda \neq 0$ the only curve on S is the image of $z = 0$. Every Hopf surface is finitely covered by a primary one.

Extending results of [GO99] Belgun proved that every Hopf surface is lcK; furthermore, the Lee form can be parallel exactly for the diagonal ones.

Inoue-Bombieri surfaces come in three different families which for simplicity we denote by S_M , S_N^- and $S_{N,u}^+$ with $u \in \mathbb{C}$; the universal covering is always a solvable group biholomorphic to $\mathbb{C} \times H$.

Tricerri [Tr82] constructed explicit lcK metrics on every surface of type S_M and S_N^- , as well on the surfaces $S_{N,u}^+$ with $u \in \mathbb{R}$. These were also the first lcK metrics with non-parallel Lee form. Belgun showed that these examples are in fact sharp.

Finally, we state the complete result

Theorem 2.1 [Be00] *(1) All diagonal Hopf surfaces and all surfaces with $Kod \geq 0$ admit lcK metrics with parallel Lee form.*

(2) All other Hopf surfaces and the Inoue-Bombieri surfaces S_M , S_N^- and $S_{N,u}^+$ with $u \in \mathbb{R}$, admit lcK metrics whose Lee form cannot be parallel.

(3) When $u \in \mathbb{C} \setminus \mathbb{R}$, the Inoue-Bombieri surfaces $S_{N,u}^+$ do not admit lcK metrics at all.

Furthermore, by Kodaira classification, every compact complex surface with b_1 odd and $\chi = 0$ belongs to exactly one of the above three classes.

In particular, contrary to the Kähler case [KS60], the family of Inoue-Bombieri surfaces $S_{N,u}^+$ shows that

Corollary 2.2 [Be00] *The class of lcK manifolds is not open under holomorphic deformations.*

3 Inoue surfaces in class VII_0^+

After the work of Belgun, Vaisman question had a complete (although not always positive) answer for all surfaces except for the class of minimal surfaces with b_1 odd and $\chi \neq 0$. We see from Kodaira classification that this in fact implies: $b_1 = 1$, $\chi = b_2 = b_2^+ > 0$ and $Kod = -\infty$. This class of surfaces is usually denoted by VII_0^+ .

The subclass with $b_2(S)$ rational curves each of which belongs to a cycle C are called Enoki [En81] and Inoue surfaces [In77]; notice that there can be at most two cycles of rational curves on a given surface [Na89, 6.3].

Enoki surfaces are characterized by the topological property that the unique cycle satisfies $C^2 = 0$; some of the Enoki surfaces also admit an elliptic curve in which case are called Parabolic Inoue surfaces. There are two more types: Hyperbolic Inoue surfaces, have two

cycles of rational curves and Half-Inoue surfaces which have a unique cycle with $C^2 < 0$; the name comes from the fact that they are double-covered by Hyperbolic Inoue surfaces.

The first examples of lcK complex surfaces in class-VII₀⁺ were found by LeBrun [Le91, p.391] on some Parabolic Inoue surfaces; this is because of the following link pointed out by Boyer, see also [Po91].

Proposition 3.1 [Bo86] *Hermitian anti-self-dual metrics on compact four-manifolds are necessarily lcK*

It was later noticed [Po97] that LeBrun's examples are in fact *bi-Hermitian*. This means that we have a four-manifold M with two integrable complex structures $J_{\pm}$ which are orthogonal w.r.t. a Riemannian metric and induce the *same orientation*. Of course we also require that there is a point $p \in M$ where $J_-(p) \neq \pm J_+(p)$; this determines the conformal structure uniquely and is equivalent to ask that the following algebraic condition holds [Tr75]

$$J_+J_- + J_-J_+ = -2a \cdot id$$

for some smooth $a : M \rightarrow [-1, 1]$ usually called the angle function [Po97]. Notice that this is a conformally-invariant notion, for which we will use the notation $(M, [g], J_{\pm})$. In order to understand the structure of such surfaces we now give a twistor presentation.

It is sometimes useful to think of an almost Hermitian structure J as a smooth section $J : M \rightarrow Z$ of the twistor space. Z is the fiber bundle of all linear complex structures at T_pM compatible with the metric and orientation; in four-dimension the fiber at $p \in M$ is the homogeneous space $SO(4)/U(2) \cong \mathbb{CP}_1$. It is known that Z is an almost complex 6-manifold - let $\mathbb{J}$ denote the almost complex structure - which only depends on the fixed conformal structure $[g]$ and orientation of M . The integrability of J is equivalent to the fact that $J : M \rightarrow Z$ is an almost $\mathbb{J}$ -holomorphic map and is also equivalent to ask that its image $J(M) =: S$ is an almost complex submanifold of $(Z, \mathbb{J})$, see for example [AMP99]; S is therefore (tautologically) biholomorphic to the original complex surface: $S \cong (M, J)$.

The twistor space Z is equipped with a natural *real structure*, namely the involution $\tau : Z \rightarrow Z$ that sends $J \mapsto -J$ and objects that are τ -invariant are then called 'real'. For example each fiber $SO(4)/U(2) \cong \mathbb{CP}_1$ is called a 'real twistor line' because it is a $\mathbb{J}$ -complex curve biholomorphic to $\mathbb{CP}_1$ and defines a non-trivial cohomology class $h \in H^2(Z, \mathbb{Z})$. It is then known that the first Chern class $c_1(Z, \mathbb{J}) = -4h$; when M is compact we also have another 'real' 4-cohomology class if we let X denote the (obviously disjoint) union $S \amalg \tau(S)$. Then its Poincaré dual X^* equals $2h$ because X intersects each real twistor line in exactly two points with the complex orientation and we can now compute the first Chern class of $(M, J) \cong S$ from adjunction formula:

$$c_1(M, J) = [c_1(Z) + X^*]_{|S} = -2h|_S \quad (1)$$

In the bi-Hermitian case there are two sections $J_{\pm} : M \rightarrow Z$; their images define two complex hypersurfaces $S_{\pm} := J_{\pm}(M)$ and we will also consider the real hypersurfaces $X_{\pm} := S_{\pm} \amalg \tau(S_{\pm})$. Notice again that there is a tautological biholomorphism $(M, J_{\pm}) \cong S_{\pm}$.

Next, we concentrate on the following intersection

$$C_+ := S_+ \cap X_- \subset Z$$

We can think of C_+ as a divisor and therefore a holomorphic line bundle on $S_+ \cong (M, J_+)$: it is a complex 1-dimensional subset because it is the intersection of (three) smooth complex hypersurfaces (with no common component) of the almost complex manifold $(Z, \mathbb{J})$. The multiplicities of the irreducible components of C_+ are the (necessarily positive) intersection multiplicities and are in any case uniquely determined by the Chern class which we compute below.

In particular, its image $C := t(C_+) \subset M$ - which is easily seen to be the set of points where $J_{\pm}$ are linearly dependent - is a closed nowhere dense subset of M [Po97, 1.3] as well as the support of the divisor C_+ .

$$C = t(S_+ \cap S_-) \amalg t(S_+ \cap \tau(S_-)) = \{p \in M \mid J_+(p) = \pm J_-(p)\}$$

The case $C = \emptyset$ is usually referred to as *strongly bi-Hermitian* because $J_{\pm}$ are linearly independent everywhere on M in this situation; see [AD08] for a complete classification.

We now proceed to compute the first Chern class of the divisor C_+ in S_+ .

Proposition 3.2 *The first Chern class of C_+ is the anti-canonical class of the surface S_+ :*

$$c_1(C_+) = -c_1(S_+)$$

i.e. on the compact complex surface (S_+) there is a holomorphic line bundle L_+ such that

- (i) $c_1(L_+) = 0$, and
- (ii) $C_+ = L_+ - K$ where K is the canonical line bundle of the surface S_+ .

Proof. In the twistor space Z we have that $C_+ \cong X_- \cap S_+$ therefore $c_1(C_+) = 2h|_{S_+}$ and the conclusion follows from (1). $\square$

Remark 3.3 *In other words C_+ is a numerically anti-canonical divisor in the complex surface S_+ . By a powerful result of Dloussky [Dl06] this forces a class- VII_0^+ surface to actually be a Kato surface. We also recall that a numerically anti-canonical divisor is automatically effective on a Kato surface [Dl06, 2.2].*

Of course we also have a numerically anti-canonical divisor C_- in the other complex surface $S_- \cong (M, J_-)$ with the same support $t(C_-) = t(C_+) = C$. This implies that $(M, J_{\pm})$ are both Kato surfaces; although they don't need to be biholomorphic in general [FP10, 7.5] the

fact that both have a numerically anti-canonical divisor with the same irreducible components and multiplicities is enough to conclude that their complex structures share a lot of common properties.

Perhaps surprisingly, the above presentation is not only useful to describe the complex structure of a possible bi-Hermitian surface but it proved to be effective in producing new examples. Starting from a Joyce twistor space [Jo95] [Fu00] over the connected sum $m\mathbb{CP}_2$ we produce in [FP10] the twistor space of bi-Hermitian anti-self-dual metrics on surfaces of class VII_0^+ with $b_2 = m$.

It is known that this kind of metrics can only leave on Parabolic or Hyperbolic Inoue surfaces and our result covers both cases; furthermore, recall that the following strong form of 3.1 applies:

Proposition 3.4 [Po97, 3.1, 3.11] *Let $(M, [g], J_\pm)$ be a compact bi-Hermitian surface. Then, the metric $[g]$ is anti-self-dual if and only if $([g], J_-)$ is lcK if and only if $([g], J_+)$ is lcK; in this situation the J_+ -Gauduchon metric is automatically J_- -Gauduchon and there are only two cases:*

- (i) *The two Lee forms coincide: $\theta_+ = \theta_-$ and $(g, J_\pm)$ span a hyper-Hermitian structure, or else*
- (ii) *The sum of the two Lee forms vanishes identically: $\theta_+ + \theta_- = 0$*

Our twistor construction produced in particular the following result:

Theorem 3.5 [FP10, 7.5, 9.3] *Every Hyperbolic Inoue surface admits anti-self-dual bi-Hermitian structures. Every Half Inoue surface admits anti-self-dual Hermitian structures. By 3.1 and 3.4 all such surfaces are lcK.*

It is also interesting to point out that bi-Hermitian metrics provide a link between lcK and generalized Kähler geometry in four-dimension. This is because a generalized Kähler structure induces a bi-Hermitian structure satisfying the second equation in 3.4, see [AD08, Prop. 3]. We can now prove the following optimal and new result about non-minimal Hyperbolic Inoue surfaces.

Theorem 3.6 *The blow up of any Hyperbolic Inoue surface at any set of points on its anti-canonical divisor admits bi-Hermitian metrics which are generalized Kähler.*

Proof. The already mentioned result [FP10, 7.5] produces in fact anti-self-dual bi-Hermitian metrics on any *properly blown-up* Hyperbolic Inoue surface - which means that the blown-up points are singular points of the anti-canonical divisor. Since the only compact hyper-Hermitian

surfaces are Hopf surfaces [Bo88], these examples must satisfy equation (ii) in 3.4 and are therefore generalized Kähler. We can now apply a recent “complementary” result of Cavalvanti-Gualtieri [CG11] which states that one can blow up smooth points of the anti-canonical divisor and still have a generalized Kähler structure. $\square$

Remark 3.7 *The condition that the blown up points belong to the anti-canonical divisor is necessary [Po97, 3.14].*

4 Kato surfaces

After the work of LeBrun and Fujiki-Pontecorvo every known example of lcK surfaces in class VII_0^+ where Inoue surfaces (a particular case of Kato surfaces) and the metrics were actually anti-self-dual and, up to double coverings, bi-Hermitian.

This situation was clarified by Marco Brunella, who left us much too early. In a sequence of two papers he proved that Kato surfaces behave extremely well with respect to the lcK property. We like to think that the results of the previous section somehow inspired him to work on the subject. This section starts with some preliminaries about Kato surfaces [Ka77].

Surfaces in class VII_0^+ all have vanishing algebraic dimension and therefore at most finitely many (holomorphic) curves. All known examples in this class are so called *Kato surfaces*: they contain a global spherical shell which means that they are obtained by holomorphic surgery from a neighborhood of the origin in $\mathbb{C}^2$ blown up at “infinitely-near” points, in particular they are diffeomorphic to $(S^1 \times S^3) \# m \overline{\mathbb{CP}}_2$ where $m = b_2(S)$.

VII_0^+ -surfaces have been intensively studied in the 1980’s by the Japanese school of Nakamura, Kato, Inoue, Enoki; they proved that every Kato surface has small deformations which are blown-up Hopf surfaces. Nakamura [Na90] also showed that every Kato surface has a singular degeneration which is a rational surface with a nodal curve; the case $b_2 = 0$ actually goes back to Kodaira [Ko64]. This characterization very much inspired our work [FP10] because we use a twistorial version of Nakamura degeneration.

There are several open problems concerning the complex structure of a surface $S \in \text{VII}_0^+$ the basic questions being: does S admit curves? Or more strongly: does it admit a cycle of rational curves? Or even more: is S necessarily a Kato surface?

Recent progress has been made by the Marseille-based group of Dloussky, Oljeklaus, Toma, Teleman; by [DOT03] a Kato surface is characterized in class VII_0^+ by having b_2 rational curves or equivalently, a numerically anti-canonical divisor [DI06]. More recently A. Teleman has shown the following result by studying the moduli space of $PU(2)$ instantons

Theorem 4.1 [Te10] *Every class VII_0^+ with $b_2 \leq 2$ has a cycle of rational curves. In particular, is diffeomorphic to $(S^1 \times S^3) \# b_2 \overline{\mathbb{CP}}_2$ and must be a Kato surface when $b_2 = 1$.*

The first results about Hermitian geometry of these surfaces were the ones of the previous sections. Thanks to the work of Brunella this situation changed dramatically soon after. His first result is that Kato surfaces are well-behaved with respect to deformations of lcK metrics.

Theorem 4.2 [Br10] *For Kato surfaces, the property of being lcK is open under holomorphic deformations.*

As a consequence, Bombieri-Inoue surfaces of type $S_{N,u}^+$ are the only known compact complex surfaces which are not lcK-stable by small deformations.

The above Theorem, combined with the results of the previous section, already gives many new lcK surfaces. However, it was greatly generalized in a following work

Theorem 4.3 [Br11] *Every Kato surface admits lcK metrics.*

This is a remarkable result, the proof is short and smart; it involves a careful construction of Kähler potentials on the modified unit ball in such a way that the holomorphic surgery becomes actually a conformal map.

To summarize, we state the results described in this paper in the following

Remark 4.4 *Theorems 2.1 and 4.3 answer Vaisman question 1.1 for all known compact complex surfaces.*

Furthermore, Nakamura strong Conjecture [Na89, 5.5] that every class VII_0^+ is actually a Kato surface, would then imply that every compact complex surface is lcK except for Bombieri-Inoue surfaces $S_{N,u}^+$ with $u \in \mathbb{C} \setminus \mathbb{R}$, which are not lcK

We finish this work with a simple observation on [Br11, question after Corollary 2].

Remark 4.5 *The universal covering is Kählerian for every known compact complex surface.*

In fact, a lcK metric is the same as a Kähler metric on the universal cover which transforms by homotheties under the action of the group of deck transformations. Furthermore, Inoue-Bombieri surfaces are covered by the product of the upper half-plane with the complex line [In74]. Therefore, the universal cover happens to be Kählerian even for the non lcK surfaces of Belgun.

References

- [AMP99] Alekseevsky D., Marchiafava S., Pontecorvo M., Compatible complex structures on almost quaternionic manifolds. *Trans. A.M.S.* **351** (1999) 997-1014.
- [AD08] Apostolov V., Dloussky G., Bihermitian metrics on Hopf surfaces. *Math. Res. Lett.* **15** (2008) 827-839.
- [AG07] Apostolov V., Gualtieri M., Generalized Kähler manifolds, commuting complex structures and split tangent bundle. *Comm. Math. Phys.* **271** (2007) 561-575.
- [AGG99] Apostolov, V., Gauduchon, P., Grantcharov, G., Bi-Hermitian structures on complex surfaces. *Proc. London Math. Soc.* (3) **79** (1999), no. 2, 414–428; *Proc. London Math. Soc.* (3) **92** (2006), no. 1, 200–202.
- [BHPV04] Barth, W., Hulek, K., Peters, C., Van de Ven, A., Compact complex surfaces, Springer, Heidelberg, Second Edition, 2004.
- [Be00] Belgun, F. A., On the metric structure of non-Kähler complex surfaces. *Math. Ann.* **317** (2000) 1-40.
- [Bg76] Bogomolov F.A., Classification of surfaces of class VII_0 and affine geometry. *Math.USSR-Izv* **10** (1976) 255-269.
- [Bm73] Bombieri, E., Letter to Kodaira, 1973.
- [Bo86] Boyer, C.P. Conformal duality and compact complex surfaces. *Math. Ann.* **274** (1986) 517-526.
- [Bo88] ———, A note on hyperhermitian four-manifolds. *Proc. Am. Math. Soc.* **102** (1988) 157-164.
- [Br10] Brunella, M. Locally conformally Kähler metrics on certain non-Kählerian surfaces. *Math. Ann.* **346** (2010) 629-639.
- [Br11] ——— Locally conformally Kähler metrics on Kato surfaces. *Nagoya Math. J* **202** (2011), 77-81.
- [Bu99] Buchdahl, N., On compact Kähler surfaces. *Ann. Inst. Fourier* **49** (1999) 287-302.
- [CG11] Cavalcanti, G. R.; Gualtieri, M., Blowing up generalized Kähler 4-manifolds. *Bull. Braz. Math. Soc. (N.S.)* **42** (2011) 537-557.

- [DI06] Dloussky, G., On surfaces of class VII_0 with numerically anticanonical divisor. *Amer. J. Math.* **128** (2006) 639–670.
- [DOT03] Dloussky, G.; Oeljeklaus, K.; Toma, M. Class VII_0 surfaces with b_2 curves. *Tohoku Math. J. (2)* **55** (2003) 283309.
- [En81] Enoki, I. Surfaces of class VII_0 with curves. *Tôhoku Math. J.* (1981) 453-492.
- [Fu00] Fujiki, A., Compact self-dual manifolds with torus actions. *J. Differential Geom.* **85** (2000), 229324
- [FP10] Fujiki, A., Pontecorvo, M., Anti-self-dual bihermitian structures on Inoue surfaces. *J. Diff. Geom.* **85** (2010) 15-71.
- [Ga84] Gauduchon P., La 1-form de torsion d’une variété hermitienne compacte. *Math. Ann* **267** (1984) 495-518.
- [GO99] Gauduchon P., Ornea L., Locally conformally Kähler metrics on Hopf surfaces. *Ann. Inst. Fourier (Grenoble)* **48** (1998) 1107-1127.
- [Gu07] Gualtieri, M., Generalized Kähler geometry. arXiv:1007.3485v1.
- [Hi06] Hitchin, N., Instantons and generalized Kähler geometry. *Commun. Math. Phys.* **265** (2006) 131-164.
- [Jo95] Joyce, D., Explicit construction of self-dual 4-manifolds. *Duke Math. J.* **77** (1995) 519552.
- [Ka77] Kato, M., Compact complex manifolds containing "global" spherical shells. *Proc. Japan Acad.* **53** (1977) 1516.
- [Ko64] Kodaira, K., On the structure of compact complex analytic surfaces. *Amer. J. Math.*, I **86** (1964) 751-798, II **88** (1966) 682-721, III **90** (1968) 55–83, IV **90** (1968) 1048-1066.
- [KM71] Kodaira, K., Morrow, J., Complex manifolds. p. 85, Holt, Rinehart & Winston, New York, 1971.
- [KS60] Kodaira, K., Spencer D.C., On deformations of complex analytic structures. III. Stability theorems for complex structures *Ann. of Math.* **71** (1960) 43-76.
- [In74] Inoue, M., On Surfaces of Class VII_0 , *Inv. Math.* **24** (1974) 269-310.

- [In77] ———, New surfaces with no meromorphic functions. II, *Complex analysis and algebraic geometry* 91106. Iwanami Shoten, Tokyo, 1977.
- [La99] Lamari, A., Courants Kählériens et surfaces compactes. *Ann. Inst. Fourier* **49** (1999) 263-285.
- [Le91] Lebrun, C.R., Anti-self-dual Hermitian metrics on blown-up Hopf surfaces. *Math. Ann* **289** (1991) 383-392.
- [LYZ94] Li, J., Yau, S.-T., Zheng, F., On projectively flat Hermitian manifolds. *Comm. Anal. Geom.* **2** (1994) 103-109.
- [Ma77] Maehara, K., On elliptic surfaces whose first Betti numbers are odd. *Proceedings of the International Symposium on Algebraic Geometry*. Kyoto Univ. (1977) 565–574.
- [Na90] Nakamura, I., On surfaces of class VII_0 with curves, II. *Tôhoku Math. J.*, **42** (1990) 475-516.
- [Na89] ——— Towards classification of nonkahlerian complex surfaces. *Sugaku Expositions, Math.* **2** (1989) 209-229.
- [Na08] ———, Survey on VII_0 surfaces. <http://www.math.sci.hokudai.ac.jp/~nakamura/70surfaces080306.pdf>
- [Po91] Pontecorvo M., Hermitian Surfaces and a Twistor Space of Algebraic Dimension 2. *Proceedings of the American Mathematical Society* **113** (1991) 177-186.
- [Po97] ———, Complex structures on four-manifolds. *Math. Ann.* **309** (1997), no. 1, 159–177.
- [Po08] ——— Bihermitian and locally conformally Kähler surfaces *Lecture note series in Mathematics, Osaka University.* **9** (2008) 172-192.
- [Sa94] Salamon S., Special structures on four-manifolds. *Riv. Mat. Univ. Parma* (4) **17** (1991) 109-123.
- [Si83] Siu, Y. T., Every $K3$ surface is Kähler. *Invent. Math.* **73** (1983) 139-150.
- [Te94] Teleman, A., Projectively flat surfaces and Bogomolov’s theorem on class VII_0 surfaces. *Internat. J. Math.* **5** (1994) 253-264.
- [Te05] ———, Donaldson theory on non-Kählerian surfaces and class VII surfaces with $b_2 = 1$. *Invent. Math.* **162** (2005) 493-521.

- [Te10] ———, Instantons and curves on class VII surfaces. *Ann. of Math.* **172** (2010) 1749-1804.
- [Th76] Thurston, W. P., Some simple examples of symplectic manifolds. *Proc. Amer. Math. Soc.* **55** (1976) 467-468.
- [Tr75] ———, Connessioni lineari e metriche hermitiane sopra varietà dotate di due strutture quasi complesse. *Riv. Mat. Univ. Parma* (4) **1** (1975) 177-186.
- [Tr82] ———, Some examples of locally conformal Kähler manifolds. *Rend. Sem. Mat. Univ. Politecn. Torino*, **40** (1982) 81-92.
- [Va80] Vaisman, I., On locally and globally conformal Kähler manifolds. *Trans. Am. Math. Soc.*, **262** (1980) 533-542.
- [Va87] ———, Non-Kähler metrics on geometric complex surfaces. *Rend. Sem. Mat. Univ. Politecn. Torino* **45** (1987) 117–123.
- [Vu09] Vuletescu, V., Blowing-up points on l.c.K. manifolds *Bulletin mathématique de la société des sciences mathématiques de Roumanie* **52** (2009) 387-393

M. Pontecorvo
 Dipartimento di Matematica
 Università Roma Tre
 L.go S.L. Murialdo 1, 00146 Roma
 max@mat.uniroma3.it